

Lena Kourkoutis and Darrell Schlom (Cornell University)

A recent study in which machine learning was applied to suggest potential high-temperature superconductors predicted that $\text{Ba}_3\text{In}_2\text{O}_6$ should superconduct at around 46 K [Z.-L. Liu *et al.* [APL Mater.](#) **8**, 061104 (2020)]. Though no barium indate superconductors are known, $\text{Ba}_3\text{In}_2\text{O}_6$ has a crystal structure that resembles $\text{YBa}_2\text{Cu}_3\text{O}_7$ and other well-known cuprate superconductors.

Thin films of $\text{Ba}_3\text{In}_2\text{O}_6$ had never been made and **PARADIM researchers were able to grow the desired phase** using a new variant of molecular-beam epitaxy (MBE), suboxide MBE, developed in PARADIM. In addition to succeeding in the growth of the targeted phase, $\text{Ba}_3\text{In}_2\text{O}_6$, **PARADIM researchers also discovered a new barium indate phase** with an even simpler crystal structure, $\text{Ba}_4\text{In}_2\text{O}_7$, that is analogous to the first high-temperature cuprate superconductor discovered, $(\text{La},\text{Ba})_2\text{CuO}_4$.

These phases are shown at the right in both cartoon form and high-resolution electron microscope images of the thin films grown. These phases are $n = 1, 2$, and ∞ members of a family of structurally related phases known as Ruddlesden–Popper phases with general formula $\text{Ba}_{n+1}\text{In}_n\text{O}_{2.5n+1}$ of which the $n=1$ member was previously unknown. The high-temperature superconducting cuprates also belong to this same family of structures. In addition to $\text{Ba}_3\text{In}_2\text{O}_6$, $\text{Ba}_4\text{In}_2\text{O}_7$ also warrants study as a potential high-temperature superconductor when doped with electrons or holes. This will be the subject of future work.

F.V.E. Hensling *et al.* [J. Vac. Science & Technol. A](#) **40**, 062707 (2022).

